International Journal of Zoology and Applied Biosciences Volume 6, Issue 4, pp: 208-211, 2021 https://doi.org/10.55126/ijzab.2021.v06.i04.017

Research Article

HYDROPHYLAX MALABARICUS A NEW DISTRIBUTIONAL RECORD FROM COIMBATORE DISTRICT, TAMIL NADU, INDIA

*1Selvaraj Selvamurugan and ²Santhana Bharathi, N.

¹Institute of Forest Genetics and Tree breeding, Coimbatore, Tamil Nadu-642 002 ²UPASITRF Tea Research Institute, Valparai, Coimbatore, Tamil Nadu

Article History: Received 11th August 2021; Accepted 22nd August 2021; Published 31st August 2021

ABSTRACT

The fungoid frog or Malabar Hills frog (*Hydrophylax malabaricus*) is a colourful frog found on the forest floor and lower vegetation. *H. malabaricus* species recorded from Nirardam, Valparai. Coimbatore district, Tamilnadu state, India. In this study reveals that the Valparaiarea is rich in amphibian diversity and support many more species. Further studies are needed on population structure, habitat use by amphibians for better understanding and also impose of several biodiversity.

Keywords: Hydrophylax malabaricus, Valparai, Frog, Malabar Hills.

INTRODUCTION

Amphibians show a very high level of diversity and endemism and are facing a global declines due to the everincreasing anthropogenic activities from the past few decades (Lawler & Mathias, 2007; Stuart et al., 2004). Among the vertebrates, amphibians are the most threatened species with extinction (IUCN, 2019). At present, India has 437 amphibian species (Frost, 2019), of these 80% are endemic and concentrated in three biodiversity hot spots, namely the Western Ghats, Himalayas and northeast India (part of Indo-Burma hot spot). About 4% of the amphibian species are critically endangered, 2% are near threatened, and others have not yet been assessed by IUCN (Dinesh et al., 2019). As more amphibian species are discovered in the country, with most of them being endemic and range restricted, it is essential to synthesize and understand the amphibian conservation in India. Fitzinger, (1843) raised genus Hydrophylax with Rana malabarica Tschudi, 1838; as the type species by original designation. However, (Günther, 1864) implied synonymy of Hydro phylax to Rana Linnaeus, 1758, and (Günther, 1864) implied synonymy of Hydrophylaxto Hylorana (an incorrect spelling of the genus Hylarana Tschudi, 1838) by placing the type species under the respective genera. (Boulenger, 1920) considered Rana malabarica as member of the

subgenus Rana. The genus Hydro phylax is diagnosed based on a combination of characters including presence of a postocular mask, robust body, rear of thighs with strong vermiculations, large rectal gland, prominent humeral gland, and ventrolateral grooves sometimes absent on finger (Oliver et al., 2015). Currently recognized species in the genus are widely distributed in Asia and are known from Sri Lanka, India, Bangladesh, southern Myanmar and western Thailand (Frost, 2004; Oliver et al., 2015). They suggested that this widespread species, which is spread across peninsular India, harbors two genetically distinct haplo groups, one restricted to the Western Ghats of Kerala and Tamil Nadu and the other distributed in the Western Ghats of Karnataka, Goa and Maharashtra and extending as far as Madhya Pradesh in central India. While the specimens from Western Ghats of Kerala and Tamil Nadu attributed to Hydrophylax malabaricus could be sensustricto (Dutta et al., 2004).

MATERIALS AND METHODS

Study area

A dam across Nirar River at a lower site, about 8 km, below the Upper Nirar weir has been constructed. This dam will serve to conduct the diverted waters of Anaimalaiyar

Basin (yet to begin) to the Sholayar valley through the unlined lower Tunnel taking off from the dam site. In addition to this, the yield of the catchment between the Upper Nirar Weir and the Dam will also be diverted to the Sholayar

Basin.In this species based on the identification by photographs. In this locality very rich biodiversity (Figure 3). Photo taken from Nirar dam valparai, Coimbatore district, Tamilnadu state, India.



Figure 1. A view of Nirar Dam, Valparai, Coimbatore district, Tamil Nadu.




Figure 2. A map showing study area of Nirar Dam, Valparai, Coimbatore district, Tamil Nadu.



Figure 3. Hydrophylax malabaricus species.

Morphology description

Vomerineteeth in two oval oblique groups between the choanae. Head moderate, depressed; snout moderate, hardly as long as the diameter of the orbit, subacuminate, moderately prominent; loreal region concave; nostril nearer to the end of the snout than to the eve; interorbital space rather narrower than the upper eyelid; tympanum very distinct, nearly as large as the eye. Fingers moderate, first extending beyond second; toes rather short, half webbed: tips of fingers and toes swollen; subarticular tubercles very strong; inner metatarsal tubercle oval, blunt; a large rounded tubercle at the base of the fourth toe; no tarsal fold. The tibio-tarsal articulation reaches the tympanum or the eye. Skin finely granulate above; a broad, not very prominent glandular lateral fold; a strong glandular fold from below the eye to the shoulder, followed by one or two glandules. Head and body bright crimson above, blackish brown on the sides; back sometimes with a few small black spots; upper lip, and a series of spots on the flank, white; limbs blackish brown above, spotted and marbled with pale brown and while; beneath uniform white, or marbled brown and white (source-Wikipedia).

RESULTS AND DISCUSSION

Although India is one of the major global biodiversity hot spots for amphibians, there is a lack of studies on conservation genetics. Globally, species that are rare, endemic, endangered, or have patchy geographic distribution have often been the target species for studying their genetic diversity so that the estimate can help in ensuring their survival either through translocation or captive breeding to avoid extinction. There is confusion regarding the type series of *Hydrophylax malabaricus*. Original description by Tschudi (1838) does not mention the number of specimens in the type series. However, (Biju et al., 2004) mentioned six specimens as syntypes of *H. malabaricus* originating from Malabar, India and collected by Roux and Dussumier. While examining the type series of *H. malabaricus* we observed that four specimens MNHN

(Biju *et al.*, 2004) designated MNHN 4440 as lectotype of *H. malabarica*. As a result, other three specimens originating from Malabar, namely MNHN 0771, MNHN 1989.3451 and MNHN 1989.3452, are paralectotypes of *H. malabaricus*. In this paper reported, Figure 3. *H. malabaricus* species, Based on the identification of photographs taken from recorded from Nirar dam, valparai, Coimbatore district, Tamilnadu state, India.

Hydrophylax bahuvistarasp. Nov. is the fourth species in the recently resurrected genus Hydrophylax, which is distributed in Sri Lanka, India, Bangladesh, southern Myanmar, and western Thailand (Frost 2015). While, H. gracilisis restricted to Sri Lanka, H. malabaricusand H. bahuvistara are currently known only from India and H. leptoglossais known from northeastern states of India, Bangladesh, Myanmar and Thailand. Within India, H. malabaricusis distributed in Kerala and Tamil Nadu, H. bahuvistara is distributed in Karnataka, Goa, Maharashtra and Madhya Pradesh, while H. leptoglossais distributed in Assam, Mizoram, Tripura, and Meghalaya. Type material of the species comes from a wide distribution. however, based on genetic data available in (Kurabayashi et al., 2005; Biju et al., 2004) and (Hasan et al., 2014) and localities for additional material from this study and distributional data for *H. malabarica* Haplo group 1 from (Biju et al., 2004), the species is widespread in peninsular India distributed in Maharashra, Karnataka, Goa and Madhya Pradesh. However, based on the current study, the populations north of Kerala belong to H. bahuvistara, which is a widely distributed species in the peninsular India, while *H. malabaricus* appears to be restricted to the Western Ghats of Kerala and Tamil Nadu. It is therefore essential to reassess the conservation status of H. malabaricus.

CONCLUSION

In India, not a single amphibian genome has been sequenced yet. As genomics can contribute immensely to the conservation of amphibians as well as provide

evolutionary insights, future studies should try to utilize genomic approaches for amphibian management. Given the population density and the ever-increasing anthropogenic pressure on the natural ecosystems in India, it is imperative that along with taxonomic and ecological studies, genetic studies should be carried out to monitor and safeguard the viability of a number of threatened amphibians. Further studies should include conservation genetics, phylogenetics and landscape genetics-based approaches, which can predict threats and risk to the amphibians, thus helping in building strategies for their conservation.

ACKNOWLEDGMENT

The authors express sincere thanks to the head of the Institute of Forest Genetics and Tree breeding, Coimbatore, Tamilnadu for the facilities provided to carry out this research work.

REFERENCES

- Biju, S. D., S. Dutta and R. Inger (2004). Hylaranamalabarica. The IUCN Red List of Threatened Species. *Version 2014.3*. Hyperlink reference not valid. Downloaded on 03 March 2015.
- Boulenger, G. A. (1920). A monograph of the American frogs of the genus Rana. Paper presented at the Proceedings of the American Academy of Arts and Sciences. 413-480.
- Dinesh, K., Radhakrishnan, C., Channakeshavamurthy, B., Deepak, P., & Kulkarni, N. (2019). A checklist of amphibians of India with IUCN conservation status. *Version* (2.0). *Online publication is available at www. zsi. gov. in. Updated till.*1-16.
- Fitzinger, L. (1843). Systema reptilium. Fasciculus primus. Amblyglossae. Braumüller et Seidel: Wien, Austria. Vindobonae, pp.106.
- Frost, D. (2004). Amphibian species of the world: an online reference, version 3.0 (22 August 2004). *Electronic*

- database accessible at http://research.amnh. org/herpetology/amphibia/index.html. American Museum of Natural History.
- Günther, A. C. L. G. (1864). *The Reptiles of British India*: Hardwicke.
- Hasan, M., Islam, M. M., Khan, M. M. R., Igawa, T., Alam, M. S., Djong, H. T., Belabut, D. M. (2014). Genetic divergences of South and Southeast Asian frogs: a case study of several taxa based on 16S ribosomal RNA gene data with notes on the generic name Fejervarya. *Turkish Journal of Zoology*, 38(4), 389-411.
- IUCN. (2019). The IUCN Redlist of threatened species (http://www. iucnredlist.org, accessed 20 January 2019).
- Kurabayashi, A., Kuramoto, M., Joshy, H., & Sumida, M. (2005). Molecular phylogeny of the ranid frogs from Southwest India based on the mitochondrial ribosomal RNA gene sequences. *Zoological Science*, 22(5), 525-534.
- Lawler, J. J., & Mathias, M. R. (2007). Climate change and the future of biodiversity in Washington: Report prepared for the Washington Biodiversity Council.1-
- Oliver, T. H., Heard, M. S., Isaac, N. J., Roy, D. B., Procter, D., Eigenbrod, F., Petchey, O. L. (2015). Biodiversity and resilience of ecosystem functions. *Trends in Ecology and Evolution*, 30(11), 673-684.
- Frost D.R. (2019). Amphibian species of the World: an online reference. *Version 6.0 accessed 10 February 2019(http://research.amnh.org/vz/herpetology/amphibi a/index.php)*.
- Stuart, S. N., Chanson, J. S., Cox, N. A., Young, B. E., Rodrigues, A. S., Fischman, D. L., & Waller, R. W. (2004). Status and trends of amphibian declines and extinctions worldwide. *Science*, 306(5702), 1783-1786.