International Journal of Zoology and Applied Biosciences Volume 8, Issue 3, pp: 27-31, 2023 https://doi.org/10.55126/ijzab.2023.v08.i03.005



Research Article

ISOLATION, ESTIMATION OF BACTERIAL LOAD, AND CHARACTERIZATION OF GUT MICROBES OF THREE GARRA SPECIES FROM BHAVANI RIVER, WESTERN GHATS, TAMIL NADU, INDIA

Yaswanthkumar S, *Raja S, Pavithra R, Kannan K, Navaneethan Muthu Krishnan

Department of Zoology, Kongunadu Arts and Science College (Autonomous), Coimbatore 641029, Tamil Nadu, India.

Article History: Received 26th March 2023; Accepted 24th April 2023; Published 10th May 2023

ABSTRACT

Microbiomes play an important role in fish physiology, particularly in digestion, by metabolizing largely indigestible feed components for the host or synthesis of essential micronutrients. The gut flora has a dynamic effect on the host's gut and systemic immunity. This study is the preliminary exploration of gut microflora of three species $Garra\ mcclellandi$, $Garra\ stenorhynchus$, and $Garra\ gotyla\ stenorhynchus$. The total bacterial load ranges from $66\times10^8\ CFU/g$ to $81\times10^8\ CFU/g$ in the gut sample. It was found that the maximum load in $G.\ stenorhynchus\ with\ 81\times10^8\ CFU/g$ followed by $Gg.\ stenorhynchus\ with\ 72\times10^8\ CFU/ml$ minimum in $G.\ mcclellandi\ with\ 66\times10^8\ CFU/ml$. The isolates were subjected to various biochemical characterizations such as the Indole test, Voges Proskauer test, Methyl Red test, Carbohydrate Utilization Test, Citrate Utilization Test, Starch Hydrolysis, and Catalase Production activity. The three fish gut samples are screened for antagonistic properties for certain human pathogens which show zero antagonistic activities. The present preliminary study shows variation in the gut microflora and these microbes may be used for the development of probiotics and drug production

Keywords: Freshwater fish, Gut microflora, Bacterial load, biochemical characterization.

INTRODUCTION

The Microbial communities are called "microbiomes" which colonize in all parts like the mucosal surfaces of fish on the skin, gills, and digestive tract (Merrifield and Rodiles, 2015). The main reason for the gut microbiome variation is genetic variation, environmental factors, or dietary habits (De Bruijn *et al.*, 2018). The exogenous enzymes which are synthesized by microbiota help to increase digestion efficiency gastrointestinal tract (Ray *et al.*, 2012). The plant starches and cellulose which are largely indigestible can be metabolized by bacterial secretions. The fish microbiomes can be enhanced productivity by increasing the feed conversion efficiency (Kuebutornye *et al.*, 2020). The mucous on the skin and gills have colonies of beneficial microbes that play an important role in fish disease resistance (Zhang *et al.*,

2018). The Gut microflora plays an important role in the digestive process, growth, and disease of the host. The gut bacterial properties such as the ability to tolerate the low pH in gastric juices resist the action of bile acids, lysozyme secreted in intestines, immune responses, and adhere to the mucus or enteric wall surface, intestinal microflora specific to each host animal and providing essential nutrients such as vitamin K (Masaaki Yoshida *et al.*, 2022). The gut microbiome produces antibacterial substances which inhibit the establishment of invading bacteria in the fish intestine (Kanagasabapathy Sivasubramanian *et al.*, 2012). The feeding strategies, developmental stages, and changing environmental conditions may change the composition of the gut microbiome (Ahmad Jammal *et al.*, 2017).

The fish in the family of Cyprinidae species are slim cyprinids with flat belly and sucking mouth (Froese et al.,

2017). Garra (sucker fish) is an omnivorous, eating alga, plankton, and small invertebrates that they suck off substrates like rocks. In this genus, intestinal length is remarkably constant within species and varies a lot between species, and it is useful to distinguish species and that dietary shifts have played a significant role in the evolution of Garra (Stiassny *et al.*, 2007). In the present study, an attempt has been made to investigate the variation in gut bacterial species in the gastrointestinal (GI) tract of freshwater fishes namely *Garra stenorhynchus*, *Garra mcclellandi*, and *Garra gotyla stenorhynchus*. The bacterial gut flora was isolated and studied the bacterial load, morphological and biochemical characterization of isolated bacteria, and analyze the diversity of bacteria based on morphological and biochemical characteristics.

MATERIALS AND METHODS

Sample collection

The fishes *Garra stenorhynchus*, *Garra mcclellandi* and *Garra gotyla stenorhynchus* were collected from the Bhavani River of western Ghats in the Coimbatore district. For each species, 15 specimens were collected and were transformed into a laboratory by using sterilized bags.

Isolation of gut microflora

The surface of the fish was sterilized with 70% alcohol and the gut was aseptically dissected from the animal's musculature under a laminar airflow chamber. The gut was weighed and placed into a 10ml of sterile double-strength phosphate-buffered saline (PBS) solution and homogenized. The bacterial suspension was streaked on nutrient agar media and the plates were incubated at 37°C for 24 h.

Spread Plate Method

A first stock solution was prepared with 9 ml distilled water in a test tube and adding 1 ml of sample to the stock solution is considered as 10⁻¹ and the same procedure is repeated for 10⁻², 10⁻³, 10⁻⁴, 10⁻⁵,10⁻⁶,10⁻⁷. The 100µl of a measured volume of each dilution is used to make a series of spread plates using an L-shape glass rod and incubated in an incubator at 37°C for 24 hours. Depending on the estimated concentration of organisms in a sample, the extent of dilution is determined. After the growth of colonies is counted using a formula

*Serial dilution = Number of colonies / Amount of plating * Dilution factor.

Morphological Characterization

Morphological Characterization of the colony was done with the help of a compound microscope. The characteristics observed include colony size, shape, color opacity, and gram staining.

Biochemical studies

The biochemical characterization of the isolated bacteria includes the Indole test, Voges Proskauer test, Methyl Red test, Carbohydrate Utilization Test, Citrate Utilization Test, Starch Hydrolysis, and Catalase Production activity.

Antagonistic Activity of Gut Isolates

The well diffusion technique was followed to determine the antagonistic property of bacteria isolated from the gut of three fishes.

RESULTS AND DISCUSSION

The total bacterial load ranges from 66*10⁸ CFU/g to 81*10⁸ CFU/g in the gut sample. It was found that the maximum load in Garra stenorhynchus was followed by Garra gotyla stenorhynchus with 72*10⁸ CFU/ml and minimum in Garra mcclellandi with 66*10⁸ CFU/ml (Figure 1). Morphological characterization techniques show the gram-positive, the microbes which retained the crystal violet were identified as gram-positive and the microbe which does not retain the crystal violet is identified as gram-negative. Among the isolates, sample 1 is positive another samples 2 and 3 are negative. Mortality of the sample 1 is negative and samples 2, and 3 were shown positive. Primary cultures were subcultures in the nutrient broth. For further studies, the single colony was isolated based on its morphology and subculture (Table 1). The biochemical tests for the isolate are given in (Table 2). The sample 1 results show that the isolates are positive for the Catalase, Voges Proskauer, Citrate utilization, and Urease, and negative for the Indole and Oxidase tests. These results confirm that the isolate is Staphylococcus species. The results of sample 2 showed that the isolates are positive for the Methyl red and Catalase test and negative for Indole, Citrate utilization test Voges Proskauer test, Urease test, and Oxidase test. These results confirm that the isolate is Salmonella species. The results of sample 3 showed that the isolates are positive for the Indole, Methyl red, and Catalase test and negative for the Citrate utilization test Voges Proskauer test, Urease test, and Oxidase test. These results confirm that the isolate is E. coli.

The antagonistic properties of a bacterial strain isolated from the gut of three fish species were (Table 3) tested for their antagonistic ability against highly virulent human bacteria such as Listeria monocytogenes, Salmonella thypi, Salmonella aureus, and E. coli. which showed negative results. The fishes have contact with the natural biota because of their aquatic environment and the various microbes are constantly ingested along with food. The transient microorganisms have a steadier and more significant collaboration with fish gastrointestinal environments (Chandni Talwar et al., 2018). Fish symbioses, co-inhabitant of fish pathogens, might be a promising source of natural antimicrobial compounds (NACS) alternative to antibiotics, limiting bacterial diseases in fishes (Rafaela et al., 2021). The analysis estimated the bacterial diversity in the gastrointestinal tract of three freshwater fishes. The bacterial load was from 66*108 CFU/g to 81*108 CFU/g in the bacterial culture. The maximum load in the sample of Garra *stenorhynchus* was followed by *Garra gotyla stenorhynchus* with 72*108 CFU/ml and the minimum load in *Garra mcclellandi* with 66*108 CFU/ml. A few variables, for example, bacterial host explicitness, food type, and water asset might make sense of these distinctions. The changes in the habitat and climate might affect the organization of the gastrointestinal miniature biota in fish (Verner-Jeffrey *et al.*, 2003).

Normally Garra benefits from the small green plants (green growth) present in the mid-water as opposed to the base mud. It doesn't uninhibitedly feed on huge plant leaves. Taking care of the environment is one of the elements that may be answerable for the presence of gut biota in these fishes. Both marine and freshwater fish have been displayed to have a particular native gut miniature biota and it might change with fish age, status, and natural circumstances (Olafsen *et al.*, 2001). Here, other than Gram-positive microbes *Staphylo coccus*, and a few Gramnegative microscopic organisms ordinarily in the gut of fish

were likewise disconnected and distinguished, like Salmonella and E. Coli. This ordinary native vegetation can seriously, barring microorganisms, restraining colonization, and, thus, preventing disease. This work firmly proposes that powerful control of microflora in fish digestion tracts is conceivable by utilizing anti-toxindelivering microbes. A comparable methodology might be conceivable in marine water fish utilizing gastrointestinal microorganisms with an inhibitory impact against pathogenic microscopic organisms. Subsequently, the current studies show the variation in the gut microflora of various fish species. This type of study will show the significance of business hydroponics as a supplement in formed fish feed or in the type of microscopic organisms' biofilm to accomplish colonization in the fish gut at a more significant level and might be helpful in the improvement of probiotics, drugs as well as in the modern chemical creation

Table 1. In this table shows the colony morphological characterization of three different samples.

Test	Sample 1	Sample 2	Sample3
Gram Staining	+	-	-
Colour	Purple	Pink	Red
Shape	Cocci	Rod-shaped	Rod-shaped
Nutrient agar	Large, Opaque, Circular,	Large, Opaque, Irregular, and	Large, Opaque, Entire,
	Smooth	Transparent	and Colorless
Motility	-	+	+

Table 2. In this table show the results of the biochemical test in three different samples (+) positive result, (-) negative result

Test	Sample 1	Sample 2	Sample3
Catalase test	+	+	+
Oxidase test	-	-	-
Indole test	-	-	+
Urease	+	-	-
Methyl red test	+	+	+
Citrate utilization test	+	-	-
Voges Proskauer test	+	-	-
Strains	Staphylococcus sp.	Salmonella sp.	Escherichia coli

Table 3. In this table shows the result of antagonistic activity of three different samples.

Name of the sample	Listeria monocytogenes	Salmonella thypi	Salmonella aureus	E. coli
Sample 1	-	-	-	
Sample 2	-	-	-	-
Sample 3	-	-	-	-

⁽⁺⁾ positive result, (-) negative result




Figure 1. This graph shows the total bacterial count in the three different samples.




Figure 2. This image shows the gram staining method's microscopic view of the three different samples. (a). Sample 1, (b). Sample 2 and, (c). Sample 3.

CONCLUSION

In this study, the preliminary exploration of gut micro flora of three species *Garra mcclellandi*, *Garra stenorhynchus* and *Garra gotyla stenorhynchus*. The total bacterial load ranges from 66×108 CFU/g to 81×108 CFU/g in the gut sample. It was found that the maximum load in *G. stenorhynchus* with 81×108 CFU/g followed by *G stenorhynchus* with 72×108 CFU/ml minimum in *G. mcclellandi* with 66×108 CFU/ml. The three fish gut samples are screened for antagonistic properties for certain human pathogens which show zero antagonistic activities.

ACKNOWLEDGMENT

The authors express sincere thanks to the Secretary, Principal, and head of the Department of Zoology, Kongunadu Arts and Science College (Autonomous), Coimbatore for the facilities provided to carry out this research work.

REFERENCES

Ahmad Jammal, (2017). Characterization of the Cultivable Gut Microflora. *Current Nutrition & Food Science*, 13, (2).

Chandni Talwar, (2018). Fish Gut Microbiome: Current Approaches and Future Perspectives, *Indian Journal of Microbiology*, 58(4), 397414. https://doi.org/10.1007/s12088-018-0760-y.

De Bruijn I, Liu Y, Wiegertjes GF, Raaijmakers JM. (2018). Exploring fish microbial communities to mitigate emerging diseases in aquaculture. *Microbial Ecology*, 94(1). doi: 10.1093/femsec/fix161. PMID: 29206925.

Froese, Rainer; Pauly, Daniel (eds.). (2017). "Garra rufa" in FishBase. August 2017 version.

Kanagasabapathy Sivasubramanian *et al.* (2012). Isolation and Characterization of Gut Micro Biota from Some

- Estuarine Fishes. *Journal of Marine Science*, 2(2), 1-6. DOI: 10.5923/j.ms.20120202.01.
- Kuebutornye, Vivian Hlordzi, Felix K.A. Gyamfua Afriyie, Emmanuel Delwin Abarike, Yishan Lu, Shuyan Chi, Melody A. Anokyewaa. (2020). The use of Bacillus species in maintenance of water quality in aquaculture: A review, *Aquaculture Reports*, (18), https://doi.org/10.1016/j.aqrep.2020.100503.
- Masaaki Yoshida. (2022). Gut microbiota analysis of Blenniidae fishes including an algae-eating fish and clear boundary formation among isolated *Vibrio* strains, *Scientific Reports*,12, 4642 https://doi.org/10.1038/s41598-022-08511-7.
- Olafsen, J.A. (2001). Interactions between fish larvae and bacteria in marine aquaculture. *Aquaculture*, 200, 223-247.
- Daniel L. Merrifield, Ana Rodiles. (2015). The fish microbiome and its interactions with mucosal tissues, Mucosal Health in *Aquaculture*, Academic Press, Pages 273-295.
- Rafaela A. Santos., Aires Oliva-Tales., Pedro Pousao-Ferreira. (2021). Isolation and Characterization of Fish gut Bacillus spp. As Source of natural Antimicrobial Compounds to fish aquaculture Bacterial Diseases, *Marine Biotechnology*. 23, 276-293.

- Ray, A.K., Ghosh, K. and Ringo, E. (2012). Enzyme producing bacteria isolated from fish gut: A review. *Aquaculture Nutrition*, doi: 10.1111/j.1365-2095.2012.00943. x.
- Stiassny, M.L.J. & Getahun, A. (2007). An overview of labeonin relationships and the phylogenetic placement of the Afro-Asian genus Garra Hamilton, 1922 (Teleostei: Cyprinidae), with the description of five new species of Garra from Ethiopia, and a key to all African species. *Zoological Journal of the Linnean Society*, 150 (1), 41-83.
- Verner-Jeffreys, D.W., Shields, R.J., Bricknell, I.R., and Birkbeck, T.H. (2003). Changes in the gut associated micro-flora during the development of Atlantic halibut (*Hippog-lossus hippoglossus L.*) larvae in three *Britishhatcheries. Aquaculture*. 219, 21-42.
- Wenjue Zhong, Yanfeng Zhang, Zihao Wu, Rongyan Yang, Xinyue Chen, Jing Yang, Lingyan Zhu. (2018). Health risk assessment of heavy metals in freshwater fish in the central and eastern North China, *Ecotoxicology and Environmental Safety*, 157, https://doi.org/10.1016/j.ecoenv.2018.03.048.