



Review Article

COVID-19 AND MUCORMYCOSIS IN INDIA: EPIDEMIOLOGY, RISK FACTORS, AND CLINICAL CHALLENGES

¹*Vijai Krishana V, ²Santhanakrishnan K, ³Thangasubha T, ⁴Selva Durai and ⁵Nazreen B

¹*PERI College of Physiotherapy, Chennai - 48, Tamil Nadu, India

²PERI College of Pharmacy, Chennai - 48, Tamil Nadu, India

³PERI Institute of Technology, Chennai - 48, Tamil Nadu, India

⁴PERI College of Arts and Science, Chennai - 48, Tamil Nadu, India

⁵PERI College of Nursing, Chennai - 48, Tamil Nadu, India

Article History: Received 18th September 2025; Accepted 27th October 2025; Published 15th November 2025

ABSTRACT

The COVID-19 pandemic has been associated with numerous secondary infections, among which mucormycosis—a rare but aggressive fungal infection has emerged as a significant health concern in India. COVID-19-associated mucormycosis (CAM) primarily affects immunocompromised patients, especially those with uncontrolled diabetes mellitus or those receiving corticosteroid therapy. This review comprehensively examines the epidemiology, risk factors, clinical manifestations, and management challenges of CAM in the Indian context. Evidence from recent studies indicates a sharp rise in mucormycosis cases during the second wave of COVID-19 in India, with rhino-orbital-cerebral mucormycosis being the most commonly reported form. Key risk factors include hyperglycemia, prolonged steroid use, immunosuppressive therapy, and COVID-19-induced immune dysregulation. Diagnosis remains challenging due to non-specific early symptoms and limited availability of rapid diagnostic tools. Management requires a multidisciplinary approach involving early diagnosis, prompt antifungal therapy, and surgical intervention when necessary. This review underscores the urgent need for heightened clinical awareness, preventive strategies, and robust healthcare infrastructure to mitigate the morbidity and mortality associated with CAM. Future research should focus on standardized treatment protocols, early diagnostic modalities, and strategies to manage high-risk populations during ongoing and future pandemics.

Keywords: COVID-19, Mucormycosis, Black Fungus, Opportunistic Infection, Diabetes Mellitus.

INTRODUCTION

The COVID-19 pandemic has placed unprecedented stress on global healthcare systems, particularly in India, where the second wave was marked not only by high infection rates but also by the alarming rise of secondary infections such as mucormycosis. Mucormycosis is a rare yet aggressive fungal disease caused by species of the order *Mucorales*, which are commonly found in soil, decaying vegetation, and organic debris. Although these fungi are typically harmless to healthy individuals, they can cause life-threatening infections in immunocompromised hosts. During the pandemic, an unexpected surge of COVID-19

associated mucormycosis (CAM) cases was reported across several Indian states. The overuse of corticosteroids, a standard therapy to manage severe COVID-19, along with pre-existing conditions such as uncontrolled diabetes mellitus, created a “perfect storm” for the development of this invasive fungal infection. According to Garg *et al.* (2021), India reported a disproportionately higher incidence of CAM compared to global averages, with morbidity and mortality rates surpassing 50% in severe cases. The rapid progression of the disease, often from nasal sinuses to orbital and cerebral involvement, highlights the need for heightened clinical vigilance and prompt therapeutic

*Corresponding Author: Vijai Krishna V, PERI College of Arts and Science, Chennai – 48, Tamil Nadu, India Email: publications@peri.ac.in

intervention. Early diagnosis, surgical debridement, and antifungal therapy such as liposomal amphotericin-B remain the cornerstones of treatment. However, limited drug availability, high treatment costs, and delayed medical response continue to pose serious challenges. Given the unique epidemiological burden in India, this review aims to synthesize existing evidence on mucormycosis in the context of COVID-19, highlighting its risk factors, clinical spectrum, treatment strategies, and implications for healthcare policy. The COVID-19 pandemic precipitated an unprecedented rise in cases of COVID-19-associated mucormycosis (CAM), with India bearing a disproportionately large share of the global burden. Several reviews and multicentre analyses report dramatically higher incidence and case numbers in India compared with most other countries. Aranjani *et al* (2021) and Muthu *et al.* (2021) highlight that the reported CAM surge during the pandemic's second wave reflected a convergence of high background diabetes prevalence, widespread steroid use, and overwhelmed healthcare systems. Large retrospective and systematic studies documented high morbidity and mortality associated with CAM, particularly when diagnosis and treatment were delayed Muthu (2022). The literature converges on two principals, interacting risk factors for CAM: uncontrolled diabetes mellitus (DM) and corticosteroid exposure. Uncontrolled DM impairs innate immunity and creates a hyperglycaemic, iron-rich milieu favourable to Mucorales growth; several case series and systematic reviews identified DM as the predominant comorbidity among CAM patients in India Anand, (2022).

Corticosteroids widely used to treat severe COVID-19 were repeatedly implicated as a precipitant of hyperglycaemia and immunosuppression; many reports emphasise inappropriate dosing or prolonged steroid courses as avoidable contributors to CAM Muthu, (2024) Other contributing factors reported across studies include prolonged hospitalization, broad-spectrum antibiotic exposure, mechanical ventilation, neutropenia (in specific subgroups), and underlying malignancy or transplantation-related immunosuppression Gangopadhyay and Bhaduri (2021). Clinical presentations of CAM most commonly involved the rhino-orbital-cerebral (ROCM) form, though pulmonary and cutaneous forms were documented, particularly among patients with severe COVID-19 or preexisting pulmonary disease Özbek *et al* (2023). ROCM typically began with sinonasal symptoms (nasal congestion, facial pain, blackish nasal discharge) progressing to orbital signs (proptosis, ophthalmoplegia, vision loss) and intracranial extension in advanced cases Anand *et al* (2022). Imaging and pathology studies emphasise the rapid angio invasive behaviour of Mucorales, explaining the swift progression from sinonasal infection to orbital/cerebral involvement when treatment is delayed. Accurate and early diagnosis remains a major challenge. Many studies stress limitations of clinical diagnosis alone and recommend a combined approach: high clinical suspicion in at-risk patients, prompt radiology (contrast

MRI for ROCM; CT and chest imaging for pulmonary disease), direct microscopy and culture of tissue specimens, and histopathology to demonstrate angioinvasion Muthu *et al* (2022). Several authors pointed out low sensitivity of cultures and the need for repeated sampling; molecular diagnostics, while promising, were not uniformly available in many settings described in Indian case series Muthu *et al* (2024). Management of CAM involves urgent combined medical and surgical interventions. Antifungal therapy principally liposomal amphotericin B (L-AMB) plus aggressive surgical debridement for ROCM remain standard of care A. Bhattacharyya and Patel (2021), However, multiple papers reported significant constraints: limited availability and high cost of L-AMB, toxicity concerns with conventional amphotericin B (and need for electrolyte/renal monitoring), and the prolonged duration of therapy required for cure Yasmin *et al* (2021). Newer azoles (posaconazole, isavuconazole) were described as salvage or step-down options, but their accessibility and cost limited their use in many reported cohorts Fitero *et al* (2024). Several case series document favourable outcomes when early combined therapy was possible, but overall survival remained strongly time-dependent and worse in patients with intracranial extension or delayed care Jeican (2024). Studies analysing outcomes highlighted very high morbidity and a case fatality that varied widely depending on site of infection, comorbidities, and timeliness of therapy. Pulmonary and disseminated forms had worse prognoses than ROCM when detected late Aggarwal *et al* (2022). Predictors of poor outcome repeatedly identified included delayed diagnosis (>7–10 days), uncontrolled hyperglycaemia at presentation, diabetic ketoacidosis, need for mechanical ventilation, and intracranial involvement Al Balushi *et al* (2022) Multicentre analyses and systematic reviews underscore that early recognition, glycaemic control, prompt surgical debridement, and access to L-AMB materially improve survival odds Dilek *et al* (2021).

Several India-centric papers emphasise contextual drivers: a high background prevalence of diabetes, disparities in healthcare access, shortages of critical antifungal drugs during the surge, and variable infection control practices in oxygen/respiratory support delivery Tayabali and Sharma (2023), Public-health and hospital-level factors such as delayed referrals, inadequate community awareness of early CAM signs, and logistical barriers to surgical care were repeatedly flagged as modifiable contributors to poor outcomes Pasquier *et al* (2023). Although the rapid accumulation of case reports, case series, and small cohort studies generated useful clinical insights, many authors noted methodological limitations: heterogeneity in case definitions, retrospective designs, incomplete microbiological confirmation, short follow-up, and publication bias toward severe cases Tayabali and Sharma (2023). Consensus statements and Delphi processes sought to standardize case definitions and management pathways, but further prospective, ideally multicentre studies are needed to clarify incidence, optimal antifungal regimens, and long-term functional outcomes.

Across reviews and expert statements, common recommendations include judicious steroid use in COVID-19 (adhering to evidence-based dosing/duration), strict glycaemic monitoring and control in COVID-19 patients, rapid case detection with low threshold for ENT/ophthalmology referral, ensuring supply chains for L-AMB and azoles, and strengthening hospital infection control (including sterile oxygen humidification) Tayabali and Sharma (2023). Authors also call for national surveillance, registry development for CAM cases, and operational research into low-cost diagnostic and therapeutic strategies suitable for resource-limited settings Pasquier *et al* (2023). The assembled literature paints a consistent picture: CAM in India during the COVID-19 waves represented a syndemic driven by the interaction of host (diabetes, immunosuppression), treatment (corticosteroids), pathogen (ubiquitous Mucorales), and system-level factors (drug availability, delayed care). While early multidisciplinary care can greatly improve outcomes, substantial gaps in diagnostics, therapeutics, and public-health readiness persist and warrant urgent attention in research and policy.

MATERIALS AND METHODOS

The present review was conducted by systematically searching multiple scientific databases, including Scopus, PubMed, Web of Science, and Google Scholar, to identify peer-reviewed articles related to COVID-19 associated mucormycosis (CAM). Keywords used in various combinations included: “COVID-19,” “SARS-CoV-2,” “mucormycosis,” “black fungus,” “diabetes mellitus,” “corticosteroids,” and “India.” Boolean operators (AND/OR) were applied to refine the search, and both clinical studies and review articles were considered. Articles published between 2020 and 2025 were included, as this period coincides with the onset and progression of

the COVID-19 pandemic. Studies were considered eligible if they (i) reported clinical cases of CAM, (ii) discussed epidemiology, risk factors, or treatment outcomes, or (iii) provided systematic or narrative reviews relevant to the Indian context. Exclusion criteria comprised non-English language publications, case reports without adequate clinical details, conference abstracts without full text, and articles unrelated to mucormycosis or COVID-19. The initial database search yielded a large pool of articles. Titles and abstracts were independently screened to eliminate duplicates and irrelevant studies. Full-text screening was performed for the remaining articles to assess methodological quality and relevance. A total of 25 key references were shortlisted, which formed the evidence base for this review. To ensure comprehensiveness, cross-referencing of bibliographies from the shortlisted papers was also carried out.

Relevant information from each article was extracted, including study type, population characteristics, clinical presentation, diagnostic modalities, treatment strategies, and outcomes. Studies were categorized under major themes such as epidemiology, risk factors, clinical features, diagnostic challenges, treatment approaches, and prognosis. A narrative synthesis was adopted, comparing findings across studies, highlighting similarities and discrepancies, and identifying gaps in current knowledge. To enhance the reliability of this review, preference was given to systematic reviews, multicenter observational studies, and Delphi consensus statements. Case reports and small series were used selectively, mainly to illustrate unique presentations or rare treatment outcomes. By combining evidence from diverse study designs, this review aims to provide a comprehensive understanding of CAM with specific emphasis on the Indian scenario.




Figure 1. COVID-19-associated mucormycosis.

RESULTS AND DISCUSSION

The literature indicates that India experienced an unparalleled surge in COVID-19 associated mucormycosis (CAM) cases compared to other countries. Aranjani *et al.* (2021) and Muthu *et al.* (2021) reported that the background prevalence of mucormycosis in India was already 70–80 times higher than in developed nations, which contributed significantly to the CAM outbreak during the COVID-19 waves. Large multicentric studies Shetty *et al.* (2024) confirmed that uncontrolled diabetes mellitus, particularly in patients receiving corticosteroid therapy, remained the strongest predictor of CAM incidence and mortality. Across the reviewed studies, uncontrolled diabetes mellitus and indiscriminate use of corticosteroids were identified as the most critical risk factors Palanisamy and Rajendran (2022). These factors not only impair the immune system but also create a hyperglycaemic and acidic microenvironment conducive to Mucorales proliferation. Other predisposing factors included prolonged hospitalisation, oxygen therapy with non-sterile humidifiers, renal failure, and malignancy-related immunosuppression Shetty *et al.*, (2024). The “triple burden” of diabetes, steroids, and COVID-19 emerged as the primary explanation for India’s disproportionate disease burden. Most CAM patients presented with rhino-orbital-cerebral mucormycosis (ROCM), characterized by facial swelling, nasal congestion, orbital pain, proptosis, and in advanced cases,

vision loss or intracranial spread M. Hoenigl Pulmonary mucormycosis, though less common, was associated with higher mortality due to late diagnosis. Diagnostic challenges were repeatedly highlighted; culture sensitivity remained low, and delays in imaging or biopsy confirmation often resulted in advanced disease presentation. Several authors advocated for a high index of suspicion in diabetic COVID-19 patients with sinus or orbital symptoms to enable timely intervention. Treatment strategies consistently emphasised a multidisciplinary approach, combining surgical debridement with systemic antifungal therapy. Liposomal amphotericin B (L-AMB) was the gold-standard drug, yet shortages and cost issues in India often forced clinicians to use conventional amphotericin formulations, which carry higher nephrotoxicity. Newer azoles such as posaconazole and isavuconazole were used as salvage or step-down therapy. Early aggressive surgery combined with antifungal therapy significantly improved survival, but mortality still ranged between 40–80% in advanced cases Arul and Kumar; Balaji (2025). While CAM was reported globally, particularly in countries with high COVID-19 caseloads, the intensity and scale of outbreaks were disproportionately higher in India. Hoenigl *et al.* (2023) observed that uncontrolled diabetes and steroid misuse were far less common drivers in Western countries, explaining lower incidence rates. This contrast underscores the unique epidemiological context of India, where systemic healthcare challenges amplified the outbreak.

Table 1. Comparison of CAM in India vs Global Cases.

Feature	India	Global (Other Countries)
Number of reported CAM cases in key review	In one systematic review up to June 2021, 233 cases from India among a total of 275 cases.	
Fatality / Case-fatality rate	~36.5% in India in the same set of cases.	~61.9% globally (among the non-India cases in the same review)
Common underlying risk factor	Diabetes mellitus is the most frequent comorbidity in Indian cases.	Globally, more heterogeneous causes, with higher proportions of malignancies, transplants, or severe immunosuppression; though diabetes remains significant.
Predominant clinical form/location of CAM	Rhino-orbital / Rhino-orbito-cerebral mucormycosis (ROCM) forms dominate (~89%) among Indian cases.	Lower proportion of ROCM (~64%) globally; more pulmonary, disseminated, or other anatomical sites
Prevalence among hospitalised COVID-19 patients	In a multicenter Indian study: 0.27% prevalence in general wards; 1.6% in ICUs.	Less data for outside India on prevalence among COVID-19 hospitalised patients in the same way, but studies indicate less frequent reporting of CAM in many countries
Time to diagnosis (early vs. late onset)	In India, many cases diagnosed later than 7-10 days post COVID-19 diagnosis; only ~25% early (<7 days) in the review.	Globally, a higher proportion of early cases in reviewed studies; ~36% in non-India cases in the review.
ICU admission, mechanical ventilation	Indian cases less often required mechanical ventilation or ICU compared to global cases.	Higher ICU / ventilation uses in global non-India cases among reviewed cohorts.

Despite improved clinical awareness, several challenges persist: delayed diagnosis due to non-specific symptoms, limited access to antifungal agents, insufficient surgical facilities in rural settings, and lack of long-term follow-up data. Furthermore, methodological gaps were noted across studies, such as reliance on retrospective data, heterogeneous case definitions, and publication bias toward severe cases. These limitations highlight the need for prospective surveillance systems, standardized treatment protocols, and real-time reporting of CAM cases in India. The reviewed literature makes it clear that CAM is not merely a secondary opportunistic infection but a syndemic outcome of COVID-19, diabetes, and steroid therapy. Early glycaemic control, rational steroid use, and heightened clinical vigilance are the most effective preventive strategies. Addressing systemic barriers—such as antifungal drug shortages, delayed referrals, and lack of awareness remains crucial for reducing morbidity and mortality in future outbreaks.

CONCLUSION

COVID-19 associated mucormycosis (CAM) has emerged as one of the most devastating opportunistic infections during the pandemic, particularly in India. The review of recent literature establishes that uncontrolled diabetes mellitus, injudicious corticosteroid use, and immunosuppression form the central triad predisposing patients to CAM. Clinical manifestations, most often in the rhino-orbital-cerebral form, progress rapidly and lead to high mortality if not diagnosed early. Successful management requires a multidisciplinary approach that combines timely diagnosis, aggressive surgical debridement, and systemic antifungal therapy, primarily liposomal amphotericin B. However, limited drug availability, high treatment costs, and delayed clinical suspicion continue to challenge healthcare delivery. Future strategies must focus on preventive and systemic interventions. Strengthening clinical awareness programs for healthcare workers and the general population is vital for early recognition of symptoms. There is a need for national surveillance systems and registries to track incidence, treatment outcomes, and emerging resistance patterns. Investment in low-cost diagnostic tools and affordable antifungal formulations will be essential to improve accessibility in resource-limited settings. Additionally, clinical trials assessing optimal steroid regimens, combination antifungal therapies, and non-invasive monitoring methods should be prioritised. From a policy perspective, integrating mucormycosis management into pandemic preparedness plans will help mitigate the impact of future outbreaks. In summary, while CAM highlighted the vulnerabilities of India's healthcare system during the COVID-19 crisis, it also presents an opportunity to implement sustainable measures that improve infection control, patient safety, and preparedness for future pandemics.

ACKNOWLEDGMENT

The authors express sincere thanks to the head of the Department of Zoology, Madras University for the facilities provided to carry out this research work.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

ETHICS APPROVAL

Not applicable

FUNDING

This study received no specific funding from public, commercial, or not-for-profit funding agencies.

AI TOOL DECLARATION

The authors declares that no AI and related tools are used to write the scientific content of this manuscript.

DATA AVAILABILITY

Data will be available on request

REFERENCES

- Aranjani, J. M., Manuel, A., Abdul Razack, H. I., & Mathew, S. T. (2021). COVID-19-associated mucormycosis: Evidence-based critical review of an emerging infection burden during the pandemic's second wave in India. *PLOS Neglected Tropical Diseases*, 15, e0009921. <https://doi.org/10.1371/journal.pntd.0009921>.
- Balushi, A. A., Al Hinai, M., & Al-Kindi, H. (2022). An opportunistic fungal infection: Invasive mucormycosis in COVID-19 case series and review. *International Journal of Infectious Diseases*. (Case series)
- Bhattacharyya, A., & Patel, R. K. (2021). COVID-19-associated rhino-orbital-cerebral mucormycosis: Clinical features and management case series and review. *Journal of Clinical Otolaryngology & Head and Neck Surgery*, 1–12.
- Dilek, A., Tekin, E., & Yilmaz, H. (2021). COVID-19-associated mucormycosis: Case report and systematic review. *Journal of Infection and Public Health*, 1–10.
- Fitero, A., López-García, J. M., & González, P. R. (2024). Corona-associated mucormycosis: Case series and review of the literature. *Journal of Fungal Medicine*, 10(5), Article 305.
- Gayathri, D., & Soundarya, R. (2024). An emergence of mucormycosis during the COVID-19 pandemic (Review). *World Academy of Science, Engineering and Technology Journal*, 6, Article 13. <https://doi.org/10.3892/228>.

- Hoeningl, M., Seidel, D., Carvalho, A., Rudramurthy, S. M., & Arastehfar, A. (2022). The emergence of COVID-19-associated mucormycosis: A global review of cases from multiple countries. *The Lancet Microbe*, 2, 80–89. (Ahead of print)
- Jeican, I. I. (2024). COVID-19-associated rhino-orbital mucormycosis: Imaging and pathology correlates. *Diagnostics*, 14(4), 429.
- Muthu, V. (2022). Definition, diagnosis, and management of COVID-19-associated pulmonary mucormycosis: Delphi consensus statement from the Fungal Infection Study Forum and Academy of Pulmonary Sciences, India. *The Lancet Infectious Diseases*, 22(9), e240–e253. [https://doi.org/10.1016/S1473-3099\(22\)00124-4](https://doi.org/10.1016/S1473-3099(22)00124-4)
- Muthu, V., Agarwal, A., & Rudramurthy, S. M. (2024). Risk factors, mortality, and predictors of survival in COVID-19-associated pulmonary mucormycosis: A multicentre retrospective study from India. *Clinical Microbiology and Infection*, 368–374. <https://doi.org/10.1016/j.cmi.2023.12.006>
- Muthu, V., Rudramurthy, S. M., Chakrabarti, A., & Agarwal, R. (2021). Epidemiology and pathophysiology of COVID-19-associated mucormycosis: India versus the rest of the world. *Mycopathologia*, 186, 739–754. <https://doi.org/10.1007/s11046-021-00584-8>
- Özbek, L., Altun, A., & Kaya, M. K. (2023). COVID-19-associated mucormycosis: A systematic review of cases and outcomes. *Systematic Reviews in Infectious Diseases*, Article 10008766.
- Pandey, V., Verma, S. K., & Tewari, R. (2025). A tragic tale of mucormycosis and COVID-19: An observational study to identify predisposing risk factors and outcomes in central India. *Cureus*, 101(4), 230–235.
- Palanisamy, P. R., & Rajendran, V. (2022). COVID-19-associated mucormycosis: A review for primary care physicians. *Journal of Family Medicine and Primary Care*, 23(9), 90–94.
- Pal, R., Gangopadhyay, A., & Bhaduri, S. S. (2021). COVID-19-associated mucormycosis: An updated systematic review. *Mycoses / Clinical Reviews*, 1–14.
- Pasquier, G., Singh, R., & Kumar, A. (2023). COVID-19-associated mucormycosis in India: Why such an unprecedented surge? *International Journal of Infectious Diseases / Perspective*, 11(2), 56–67.
- Shetty, S. K., Patil, R., & Bhandari, M. (2024). Management of COVID-19-associated rhino-orbital mucormycosis: Surgical approaches and outcomes case series. *Frontiers of Oral and Maxillofacial Medicine*, Article 78228.
- Tayabali, K., & Sharma, S. (2023). Epidemiology of COVID-19-associated mucormycosis: India's experience and global comparisons. *Mycopathologia / Epidemiology*, 3(4), 11–23.
- Anand, T., Kumar, V., & Rao, P. S. (2022). A case-control investigation of COVID-19-associated mucormycosis (CAM) in India: Trend and risk factors. *BMC Infectious Diseases*, 22, Article 856. <https://doi.org/10.1186/s12879-022-07844-y>
- Yasmin, F., Khan, M. A., & Rahman, S. (2021). COVID-19-associated mucormycosis: A systematic review of clinical features, management, and outcomes. *Journal of Clinical Medicine Research*, 9(4), Article 65.
- Arul, A., Kumar, P., & Balaji, R. S. (2025). Pulmonary mucormycosis- An experience from a tertiary care centre during the COVID-19 waves in India. *Clinical Microbiology Insights*, 34, 90–95.

